

University of Groningen

Annulation reactions of hetero-aromatic compounds. A new synthetic route to benzo[b]thiophene, benzo[c]thiophene and quinoline derivatives

Terpstra, Jan Willem

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1982

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Terpstra, J. W. (1982). *Annulation reactions of hetero-aromatic compounds. A new synthetic route to benzo[b]thiophene, benzo[c]thiophene and quinoline derivatives*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

S U M M A R Y

In this thesis the annulation reactions of *ortho*-disubstituted hetero-aromatic compounds with Michael acceptors are described.

In Chapter I a general introduction is given to annulation reactions. The choice of the hetero-aromatic compounds that we used for our annulations, *i.e.* thiophenes and pyridines, is elucidated. Furthermore, properties of thiophenes and pyridines with respect to their substitution and annulation reactions are discussed. Existing annulation routes to benzo[*b*]thiophenes, benzo[*a*]thiophenes and quinolines, and substitution reactions of these groups of compounds, also get attention in Chapter I.

Chapter II describes the synthesis of the starting materials for the annulation reactions, that is:



$n = 1 \text{ or } 2$

$E = \text{COOMe, CN or CHO}$

$n = 1 \text{ or } 2$



$E = \text{COOMe or CN}$

$n = 1 \text{ or } 2$

$E = \text{COOMe}$

The general requirements, *i.e.* a nucleophilic and an electrophilic centre, both α to the heterocycle and in *ortho*-positions, and the specific choice of substituents for our starting materials are discussed. These compounds are all new compounds. Comments are given on the problems encountered in the synthesis of 3,4-disubstituted pyridines. We have been unable to solve these problems.

The annulation reactions of 2,3-disubstituted thiophenes to benzo[*b*]thiophenes and 4,5-dihydrobenzo[*b*]thiophenes are described in Chapter III.



Various reaction conditions for these annulation reactions were investigated. In this chapter X-ray proof of the (dihydro)benzo[*b*]thiophene struc-

ture and of the s
as ^1H and ^{13}C NMR
reactions is also
to convert 4,5-di
Chapter IV dis
phenes with Micha
derivatives.

^1H and ^{13}C NMR s
port the benzo[*a*]
annulation react
critically evalu
In the last C
pyridines with M
quinolines.

^1H and ^{13}C N
A comparison of
V showed the fo
disubstituted p
thiophenes.

ture and of the stereochemistry of the annulation products is given, as well as ^1H and ^{13}C NMR data of these products. The mechanism of the annulation reactions is also discussed in Chapter III. Further, reactions are described to convert 4,5-dihydrobenzo[*b*]thiophene products into benzo[*b*]thiophenes.

Chapter IV discusses annulation reactions of 3,4-disubstituted thiophenes with Michael acceptors. These reactions lead to benzo[*c*]thiophene derivatives.



^1H and ^{13}C NMR spectral data of the products are analyzed and used to support the benzo[*c*]thiophene structure. The analogies and differences in the annulation reactions to benzo[*c*]thiophenes and to benzo[*b*]thiophenes are critically evaluated.

In the last Chapter, V, reactions are discussed of 2,3-disubstituted pyridines with Michael acceptors, leading to quinolines and 7,8-dihydroquinolines.



^1H and ^{13}C NMR data supported the quinoline structure of the products. A comparison of the annulation reactions described in Chapter III, IV, and V showed the following reactivity order towards Michael acceptors: 2,3-disubstituted pyridines > 2,3-disubstituted thiophenes > 3,4-disubstituted thiophenes.